Draft version December 2, 2011
Preprint typeset using LaTeX style emulateapj v. 11/10/09

arXiv:1112.0179v1 [astro-ph.GA] 1 Dec 2011

THE INNER GALACTIC BULGE: EVIDENCE FOR A NUCLEAR BAR?

Ortwin Gerhard[1] and Inma Martinez-Valpuesta[1]
Max-Planck-Institut für Extraterrestrische Physik, Giessenbachstrasse, 85748 Garching, Germany

Draft version December 2, 2011

ABSTRACT

Recent data from the VVV survey have strengthened evidence for a structural change in the Galactic bulge inwards of $|l| \leq 4°$. Here we show with an N-body barred galaxy simulation that a boxy bulge formed through the bar and buckling instabilities effortlessly matches measured bulge longitude profiles for red clump stars. The same simulation snapshot was earlier used to clarify the apparent boxy bulge - long bar dichotomy, for the same orientation and scaling. The change in the slope of the model longitude profiles in the inner few degrees is caused by a transition from highly elongated to more nearly axisymmetric isodensity contours in the inner boxy bulge. This transition is confined to a few degrees from the Galactic plane, thus the change of slope is predicted to disappear at higher Galactic latitudes. We also show that the nuclear star count map derived from this simulation snapshot displays a longitudinal asymmetry similar to that observed in the 2MASS data, but is less flattened to the Galactic plane than the 2MASS map. These results support the interpretation that the Galactic bulge originated from disk evolution, and question the evidence advanced from star count data for the existence of a secondary nuclear bar in the Milky Way.

Subject headings: Galaxy: structure — Galaxy: bulge — Galaxy: evolution — methods: numerical

1. INTRODUCTION

The barred nature of the Milky Way is well-established from NIR photometry (Blitz & Spergel 1991; Dwek et al. 1995; Binney et al. 1997; Bissantz & Gerhard 2002) and star counts (Stanek et al. 1994; López-Corredoira et al. 2005; Benjamin et al. 2005; Cabrera-Lavers et al. 2007), from comparing HI and CO lv-diagrams with hydrodynamic models (Englmaier & Gerhard 1999; Fux 1999; Bissantz et al. 2003), and from dynamical modelling of the kinematics of bulge stars (Zhao 1996; Fux 1997; Bissantz et al. 2004; Shen et al. 2010).

One open issue is the structure of the inner Galactic bulge, for $R \lesssim 0.5$ kpc. Is this simply the scaled-down, higher-density parts of the surrounding boxy bulge? Does it become axisymmetric towards the center? Does it contain a "classical" bulge component? Or does it even contain a secondary nuclear bar as do about one third of barred galaxies like ours (Erwin 2011)?

This issue is important; for example, a nuclear bar misaligned with the main Galactic boxy-bulge bar would substantially modify the morphology of the gas flow in the central Galaxy, as illustrated by the simulations of Rodriguez-Fernandez & Combes (2008, hereafter RFC08) and Namekata et al. (2009).

Nishiyama et al. (2005) analyzed red clump (RC) star counts in the inner bulge. Because RC stars are approximate standard candles, their magnitude distribution is a proxy for the stellar density along the line-of-sight (Stanek et al. 1994). Determining the maximum of the RC magnitude distribution at various longitudes in a strip at $b = -1°$, Nishiyama et al. (2005) found a clear change of slope in the RC longitude profile at $|l| = 4°$, separating a steeper slope in the outer bulge from a shallow slope in the nuclear regions.

Recently, Gonzalez et al. (2011a) analyzed new RC star counts from the VVV survey and found excellent agreement with Nishiyama et al. (2005). They also determined the RC maximum versus longitude for $b = +1°$ and found good agreement with the $b = -1°$ results, as expected for a triaxially symmetric structure. This also shows that reddening corrections, which differ greatly between these latitudes, are unlikely to cause the observed change of slope in the longitude profiles.

The RC star counts therefore indicate a significant and fairly sudden change in the structure of the bulge at $|l| \simeq 4°$. Here we analyze the distribution of RC maxima for a suitably oriented boxy bulge - barred galaxy model. Surprisingly, we find that the RC longitude profiles predicted by this model are very similar to the observed profiles. We show that this can be traced back to the model's density structure becoming more axisymmetric towards the center, and finally we compare the asymmetry of the predicted star count map in the inner bulge region with available analysis of 2MASS data.

2. A MILKY WAY-LIKE N-BODY MODEL

The model analyzed in this paper stems from the barred galaxy simulation of Martinez-Valpuesta et al. (2006) and is the same model which we used earlier in order to argue that the long bar and the boxy bulge in the MW are the three-dimensional and planar components of the MW's main bar (Martinez-Valpuesta & Gerhard 2011, hereafter MVG11). It evolved from an initially exponential disk with $Q = 1.5$ embedded in a dark matter halo, and developed a prominent boxy bulge through a buckling instability after ~ 1.5 Gyr. To resolve the nuclear bulge better the particle distribution is symmetrized with respect to the midplane. We consider the simulated galaxy at time ~ 1.9 Gyr, some time after the instability when the bar has resumed its evolution and has regrown through further angular momentum transfer to the halo. The snapshot chosen for our analysis in this paper is that shown in Fig. 1 of MVG11; however, we have checked that the results given for the nuclear regions below do not significantly change when the snap-



FIG. 1.— Distributions of simulated red clump distance moduli in different bulge fields (black lines), for an N-body model of the Galactic boxy bulge and bar. The model is viewed from a position $R_0 = 8$kpc from the Galactic center and $\alpha_{\rm bar} = 25°$ away from the bar major axis. The particle positions are convolved with a Gaussian luminosity function for the red clump stars with width $\sigma_{\rm RC} = 0.17$ mag. Gaussian fits to the distributions are shown by dotted red lines.

shot in Fig. 2 of MVG11 is used instead.



FIG. 2.— Maxima of observed and model magnitude distributions for red clump (RC) giant stars in bulge fields as a function of longitude. Top: Simulated RC maxima for strips with latitudes $b = [1 \pm 0.5°]$ and $[0 \pm 0.5°]$ (black dots), compared with data from the VVV survey at $b = \pm 1°$ (Gonzalez et al. 2011a, open squares). We used $M_{\rm K} = -1.70$ to shift the model distance moduli to the magnitude scale of the data; see text. Bottom: RC maxima in strips with different latitudes, colour-coded as indicated on the plot. Individual fields have sizes $\Delta l = \Delta b = 1°$ and are separated in longitude by 1°. The change of slope in the inner few degrees seen at low latitudes is absent at $b = 5°$. The horizontal line illustrates the assumed distance to the GC.

2.1. *Magnitude-dependent simulated star counts in the inner bulge*

The stellar population of the Galactic bulge is predominantly old ($\sim$ 10 Gyr) and has a metallicity distribution with FWHM $\sim$ 1 dex around [Fe/H]$\simeq$-0.1 dex (Brown et al. 2010). In this range the number of RC stars per unit mass varies within $\sim$ 10% (Salaris & Girardi 2002), so we can assume that the spatial distribution of RC stars is the same as that of model particles (mass). We project the bulge of this model onto the sky in longitude-latitude coordinates (l, b) as seen by an observer at $R_0 = 8$ kpc distance from the Galactic center (GC hereafter), whose line-of-sight (LOS) is at an angle $\alpha_{\rm bar} = 25°$ relative to the long axis of the bar (Gerhard 2002, MVG11). Dividing the projected particle distribution into fields of $\Delta l = \Delta b = 1°$, we sort the particles in each field by distance modulus. To account for the finite width of the luminosity function for red clump giant stars (RC hereafter), we convolve each particle with a Gaussian of width $\sigma = 0.17$ mag (Alves 2000). The absolute M_K(RC) varies by $\sim$ 0.26 mag over the observed FWHM metallicity range (Salaris & Girardi 2002), corresponding to a σ of 0.11 mag. Added in quadrature, this gives a total $\sigma' = 0.21$ mag. Using σ' instead of $\sigma = 0.17$ mag has a negligible effect on the figures shown below.

Fig. 1 shows the resulting LOS magnitude distributions for fields with $b = 0$ and several l. The main peak corresponds to the inclined bar and boxy bulge; it shifts from m-M$\simeq$ 14.28 at $l = 10°$ on the bar's near side to m-M$\simeq$ 15.14 at $l = -10°$ on the far side. The background disk is seen clearly at m-M$\simeq$ 15.55 in the top panel for $l = 10°$; the slope towards m-M=16 is consistent with the disk scale-length. This disk feature is similar in absolute numbers but weaker in relative numbers for smaller $|l|$ and seems to blend with the far end of the bar at $l = -10°$. A similar phenomenology appears to be seen in the VVV data (Gonzalez et al. 2011a, Fig. 2), which in addition show a 'background' of K giants not included in the Gaussian RC luminosity function.

We determine the maxima of the simulated RC distribution in all fields by fitting a Gaussian to the brightest peak of the distribution. Fig. 2 shows the resulting RC longitude profiles for several latitude strips, and compares those at $b = 0°$ and $b = 1°$ to the results from Gonzalez et al. (2011a) at $b = \pm 1°$. This last step requires choosing an absolute K magnitude for the RC. To shift the model to the magnitude scale of the data, we need to use M_K(RC)= -1.70. The most recent M_K(RC)=-1.61 (Alves 2000; Laney et al. 2011) for the local RC, corrected for population effects (Salaris & Girardi 2002), would predict M_K(RC)$\simeq -1.52$. The -1.70 value would thus argue for a shorter distance R_0 to the GC, while the best current value is $R_0 = 8.3 \pm 0.23$ kpc (Brunthaler et al. 2011). We do not



FIG. 3.— *Left panel*: Face-on surface density of the particles with $|z| < 300$ pc, with overplotted maxima of the line-of-sight density distributions (open circles) and maxima of the simulated line-of-sight RC magnitude distributions (full circles) for particles in the latitude range $|b| \leq 2°$. *Right panel*: Same for the particles with 450pc $< |z| <$ 750 pc, and the latitude range $3° < b < 5°$. The long axis of the bar is at an angle $\alpha_{\rm bar} = 25°$ relative to the line-of sight to the observer at the assumed galactocentric radius $R_0 = 8$ kpc. Dotted lines show directions $l = 0, \pm2, \pm4, \pm6, \pm8, \pm10°$, as seen from the observer's position.

pursue this issue further here.

Both the observed $b = \pm1°$ longitude profiles and the low-latitude model longitude profiles show the same qualitative result: the steep slope of the RC maxima seen at large positive and negative $|l| > 4°$ flattens for small $|l|$[1]. The model profile is slightly flatter than the data inside $l = \pm2°$ and at $l > 4°$, but steeper around $l = -4°$. The origin of the difference at $l > 6°$ between model and data is unclear; the maxima in the observed distributions are very broad and the contribution from the disk substantial, but also differential population effects might contribute. By contrast with the lower latitudes, the $b = 5°$ model longitude profile shows a markedly weaker flattening in the inner parts $|l| < 4°$.

This model was not constructed to match Milky Way observations – it is a generic boxy bulge and bar model from a simulation; thus we do not expect a perfect match with any Milky Way data. The agreement between the low-latitude observed and model profiles is therefore even more striking. Previously, the flattening of the observed longitude profiles has been interpreted as due to a distinct structure in the inner Galactic bulge (Nishiyama et al. 2005; Gonzalez et al. 2011a), perhaps a nuclear bar (see also Alard 2001, RFC08). The simulated model bulge does not contain such a structure.

2.2. *Origin of the change of slope in the simulated star counts*

To understand the cause of the flattening in the model longitude profiles, let us first consider a simple thought experiment. We know that the RC maxima for a thin bar follow approximately the major axis of the bar (e.g. Cabrera-Lavers et al. 2007). For a thicker bar, e.g., with planar axis ratio $b/a \sim 0.5$, a volume effect arises due to the geometry of the cones with fixed $\Delta l, \Delta b$. This causes the maxima of the RC magnitude distributions to lie behind the maxima of the line-of-sight (LOS) density distributions (Cabrera-Lavers et al. 2007, MVG11). However, as long as the density distribution is approximately scale-free, we would expect the RC longitude profile to have a similar slope as for the thin bar, following the orientation of the major axis.

Now insert an axisymmetric component with high stellar density in the center of the bar. If its density is sufficiently high, it will dominate the surrounding bar, and since it is axisymmetric, the slope of its longitude profile will be nearly zero. By lowering the density of the central component, we will therefore be able to arrange a slope that is between zero and that of the surrounding barred bulge. This simple reasoning shows that the result of the previous subsection may be explained if the center of the barred bulge has sufficient central concentration and is rounder than the outer bar. While if the shape of the density distribution were to remain independent of radius and only the central concentration were increased, in first order only the total number of stars in the central LOS would be changed, which does not shift the maxima in the histograms.

Fig. 3 shows the surface density of the barred model galaxy in face-on projection, for two slices through the model. We see from the left panel that the near-planar part of the boxy bulge-bar shows a significant change in the axis ratio with major axis length, such that the central parts are nearly round (ellipticity $\epsilon \sim 0.8$) while the outer bar is much more elongated ($\epsilon \sim 0.4$). At the same time, from the spacing of the contours we see a change of slope in the major axis density profile, which sets in at approximately major axis length $\simeq 700$ pc corresponding in projection to $l \simeq 4°$. The maxima of the LOS density distributions and the RC magnitude distributions overplotted on the surface density clearly show the resulting change of slope in the inner few degrees. Note also that the steep outer slope on the $l > 0$ side is significantly decreased by the volume effect, and that on the $l < 0$ side the outer slope is increased.

---

[1] Due to the symmetrization the model star counts at $b = 1°$ and $b = -1°$ are equivalent.

The Gaussian dispersions found from the longitude profile fits in Fig. 1 at $l = \{-10°, -4°, -1°, 1°, 4°, 10°\}$ are $\{0.28, 0.31, 0.21, 0.20, 0.24, 0.43\}$ at $b = 0°$ (Fig. 1) and $\{0.33, 0.29, 0.21, 0.20, 0.24, 0.40\}$ at $b = 1°$, compared to the average VVV values at similar longitudes and $b = \pm 1°$ of $\{0.41, 0.33, 0.28, 0.30, 0.33, 0.44\}$. Averaged over the model longitude profiles for $|l| \leq 4°$ and $b = \{0°, 1°\}$, the average Gaussian $\sigma_{\pm 4} \simeq 0.24$ mag, corresponding to a deconvolved FWHM of 0.40 mag or a FWHM in the LOS density distribution of ±750 pc. To compare with the VVV data, we need to convolve with the stellar population broadening $\sigma_{\rm popn} \simeq 0.11$ mag from Section 2.1, and with the broadening by residual patchy extinction effects within the 3'-6' subfields used for deriving the reddening map (Gonzalez et al. 2011b). An upper limit for the latter at $b = \pm 1°$ can be estimated by comparing the width in J-K colour of the dereddened RC in a field with $\rm A_K = 1.5$ with a nearby low-extinction field at $b \simeq -1$. The observed difference results in $\sigma_{\rm J-K} = 0.28$ mag (O. Gonzalez, private communication), from which $\sigma_{\rm K} \simeq 0.6\sigma_{\rm J-K} = 0.17$ mag. Adding both values in quadrature to the model $\sigma_{\pm 4} \simeq 0.24$ mag predicts a total $\sigma = 0.31$ mag, which is close to the observed width for the VVV data, $\sigma_{\pm 4} \simeq 0.30$ mag. The intrinsic RC widths of the model and of the Galactic bulge as traced by the VVV RC data may therefore be quite similar. For comparison, the projected $\Delta l = \pm 4° \simeq \pm 560$pc. Therefore, for both model and data the structure inside $l = \pm 4°$ is more extended along the LOS than in l.

By contrast, for the slice with $450\text{pc} < |z| < 750$ pc shown in the right panel of Fig. 3 the change of axis ratio in the barred bulge is much less pronounced, and also the change of the major axis density slope is less pronounced. As a result, the density and RC maxima at higher latitudes lie on more nearly straight lines.

Barred galaxies are often characterized by a rising ellipticity profile at constant position angle, up to a maximum (e.g. Peletier et al. 1999), for both disk- and bulge dominated galaxies (Barazza et al. 2008). N-body models of bars have a range of morphologies; but isodensities have not been studied extensively. Some models in Fux (1997); Athanassoula & Misiriotis (2002) appear similar to the N-body model used above.

2.3. *Total star count map in the inner bulge*

2MASS star count maps for the Galactic bulge in the inner few degrees show a longitudinal asymmetry (Alard 2001, RFC08) which these authors interpreted as independent evidence for the possible existence of a separate nuclear bar. How does the nuclear bulge of our simulation compare to these data? Figure 4 shows the smoothed model star count map in 13° × 8° and in the central 3° × 2°. To construct this map, all particles in the simulation have been used, i.e., possible incompletenesses in the data at the far side of the bulge are not modelled. We see that the apparent highest surface density is shifted to $l \simeq -0.3°$, as are the centroids of the surrounding contours. Also the vertical extent of the isodensities is slightly larger on the $l < 0°$ side for small $|l|$, but becomes larger on the $l > 0°$ side for larger $|l|$. These asymmetries resemble those predicted by Blitz & Spergel (1991) for the integrated light from the central region of an inclined triaxial bulge; however, the star counts here are weighted by two powers of distance more than the integrated light.



Fig. 4.— Map of the particle surface density in a longitude-latitude plot as seen by an observer 8 kpc from the model galaxy's center. Top: central few degrees; bottom: inner bulge region. Contours are logarithmically spaced. The bar angle $\alpha_{\rm bar} = 25°$.

The published 2MASS star count maps (Alard 2001, RFC08) qualitatively show the same asymmetry pattern. The most significant difference to the model is that the innermost contours in the observed maps are significantly more flattened than the corresponding model contours. These innermost 2MASS contours are possibly affected both by incompleteness (which however would probably decrease the apparent flattening) and by blending (which might increase it). Similar maps for the VVV data would therefore be very valuable. Also, the model used here has a somewhat thicker disk than the Milky Way and, most importantly, it does not include any dissipative evolution such as gas infall and star formation on the orbits around the ILR, while such star formation is clearly on-going in inner Galaxy (Yusef-Zadeh et al. 2009). Understanding whether the flattened component seen in the central 2MASS star counts is consistent with a symmetric disk-like nuclear bulge (Launhardt et al. 2002) or is related to a nuclear bar will require much more detailed data and analysis. Here we merely emphasize that the asymmetry structure seen in the 2MASS data is not a tell-tale evidence for such a secondary bar.

3. DISCUSSION AND CONCLUSIONS

We have shown that the RC maximum longitude profiles measured for $b = \pm 1°$ are well-matched by an N-body model of a boxy bulge and bar which formed from a bar-unstable disk. This model is the same as in MVG11, and we have used the same orientation and scaling to avoid parameter fitting. The change in slope of the model profiles in the inner few degrees is caused by a transition from highly elongated to nearly axisymmetric density contours around this scale. The average FWHM of the simulated RC profiles in the range $l = \pm 4^\circ$ ($\simeq \pm 560$ pc at $R_0 = 8$ kpc) is $\simeq 0.56$ mag, consistent with the VVV data after convolution with estimated stellar population and patchy extinction broadening. This corresponds to a deconvolved bulge FWHM scale of ±750 pc along the LOS in this region. At higher latitudes the transition in the model isodensities weakens, predicting that the change of slope in the longitude profile disappears.

The projected star count map of the model has the same asymmetry pattern as inferred previously from 2MASS observations. The central surface density contours in the Galactic bulge are significantly more flattened, however. This could imply the existence of a dense central disk around the location of the ILR.

Our results generally support the idea that the main part of the Galactic bulge formed from the disk through dynamical instabilities. They also show that the evidence from star counts so far does not require a secondary nuclear bar in the Milky Way.

ACKNOWLEDGMENTS

We thank R. Benjamin, O. Gonzalez, D. Minniti, and M. Rejkuba for helpful discussions on asymmetries in the GLIMPSE and VVV surveys, the anonymous referee, and the Aspen Center for Physics for their hospitality when this work was started. This paper is based on work supported in parts by the SPP 1177 program of the German Science Foundation under Grant GE 567/4-2 and by the US National Science Foundation under Grant No. 1066293.

REFERENCES

Alard, C. 2001, A&A, 379, L44
Alves, D. R. 2000, ApJ, 539, 732
Athanassoula, E., & Misiriotis, A. 2002, MNRAS, 330, 35
Barazza, F. D., Jogee, S., & Marinova, I. 2008, ApJ, 675, 1194
Benjamin, R. A., Churchwell, E., Babler, B. L., Indebetouw, R., & et al. 2005, ApJ, 630, L149
Binney, J., Gerhard, O., & Spergel, D. 1997, MNRAS, 288, 365
Bissantz, N., Debattista, V. P., & Gerhard, O. 2004, ApJ, 601, L155
Bissantz, N., Englmaier, P., & Gerhard, O. 2003, MNRAS, 340, 949
Bissantz, N., & Gerhard, O. 2002, MNRAS, 330, 591
Blitz, L., & Spergel, D. N. 1991, ApJ, 379, 631
Brown, T. M., et al. 2010, ApJ, 725, L19
Brunthaler, A., Reid, M. J., Menten, K. M., Zheng, X.-W., & et al. 2011, Astronomische Nachrichten, 332, 461
Cabrera-Lavers, A., Hammersley, P. L., González-Fernández, C., López-Corredoira, M., & et al. 2007, A&A, 465, 825
Dwek, E., Arendt, R. G., Hauser, M. G., Kelsall, T., & et al. 1995, ApJ, 445, 716
Englmaier, P., & Gerhard, O. 1999, MNRAS, 304, 512
Erwin, P. 2011, Memorie della Societa Astronomica Italiana Supplementi, 18, 145
Fux, R. 1997, A&A, 327, 983
—. 1999, A&A, 345, 787
Gerhard, O. 2002, in Astronomical Society of the Pacific Conference Series, Vol. 273, The Dynamics, Structure & History of Galaxies, ed. G. S. Da Costa & H. Jerjen, 73–+
Gonzalez, O. A., Rejkuba, M., Minniti, D., Zoccali, M., Valenti, E., & Saito, R. K. 2011a, A&A, 534, L14
Gonzalez, O. A., Rejkuba, M., Zoccali, M., Valenti, E., & Minniti, D. 2011b, A&A, 534, A3
Laney, C. D., Joner, M. D., & Pietrzynski, G. 2011, ArXiv e-prints
Launhardt, R., Zylka, R., & Mezger, P. G. 2002, A&A, 384, 112
López-Corredoira, M., Cabrera-Lavers, A., & Gerhard, O. E. 2005, A&A, 439, 107
Martinez-Valpuesta, I., & Gerhard, O. 2011, ApJ, 734, L20+
Martinez-Valpuesta, I., Shlosman, I., & Heller, C. 2006, ApJ, 637, 214
Namekata, D., Habe, A., Matsui, H., & Saitoh, T. R. 2009, ApJ, 691, 1525
Nishiyama, S., Nagata, T., Baba, D., Haba, Y., & et al. 2005, ApJ, 621, L105
Peletier, R. F., Knapen, J. H., Shlosman, I., Pérez-Ramírez, D., Nadeau, D., Doyon, R., Rodriguez Espinosa, J. M., & Pérez García, A. M. 1999, ApJS, 125, 363
Rodriguez-Fernandez, N. J., & Combes, F. 2008, A&A, 489, 115
Salaris, M., & Girardi, L. 2002, MNRAS, 337, 332
Shen, J., Rich, R. M., Kormendy, J., Howard, C. D., De Propris, R., & Kunder, A. 2010, ApJ, 720, L72
Stanek, K. Z., Mateo, M., Udalski, A., Szymanski, M., Kaluzny, J., & Kubiak, M. 1994, ApJ, 429, L73
Yusef-Zadeh, F., et al. 2009, ApJ, 702, 178
Zhao, H. 1996, MNRAS, 283, 149